

Dairy-Mel products are formulated for utilisation with almost any roughage source for high milk production as well as high milk solid outputs.

- Dairy-Mel products contain high levels of bypass proteins which are highly digestible to increase milk production and optimize milk protein levels.
- Dairy-Mel products contain at least 35% maize.

- Dairy-Mel products contain ionophores and other medicaments for improved rumen health and feed conversion.
- Dairy-Mel products are palatable.

COMPOSITION	Dairy-Mel Energy Balancer	Dairy-Mel 17	Dairy-Mel 19	Dairy-Mel 21
Class:	Concentrated Dairy Feed			
Registration Number:	V23292	V22384	V22381	V22383
AFGRI Product Code:	R1028	R1024	R1025	R1026
Nutrients	g/kg	g/kg	g/kg	g/kg
Crude Protein (min.)	150	170	190	210
Crude Protein from NPN (max.)	19.13 %	16.88 %	15.11 %	13.67 %
Urea (max.)	10	10	10	10
Moisture (max.)	120	120	120	120
Crude Fat (min.)	25	25	25	25
Crude Fat (max.)	100	85	85	85
Crude Fibre (max.)	120	120	120	120
Calcium (min.)	8	8	8	8
Calcium (max.)	13	13	13	13
Phosphorus (min.)	4	4	4	4

FEEDING RECOMMENDATIONS

The quantity of dairy feed fed per animal will depend on:

- 1. The quality and availability of grazing and/or roughage.
- 2. The stage of lactation.
- 3. The milk and butterfat production.

Form: Mash or Pellets

Packages: Bulk or 40 kg Bags

WARNING

This farm feed contains a NPN source and registered stock remedies. Feed strictly according to instructions.

FEEDING INSTRUCTIONS

- Adaptation of animals is important when changing from a NPN-free dairy meal to a dairy meal which contains NPN (eg. urea). It is advisable to use a 50/50 mixture of the two types of meal over a period of 4-6 days in order to facilitate adaption.
- It is recommended that the dairy meal be fed in at least 2-4 meals per day or as a TMR (Total Mixed Ration).
- This dairy meal should not be fed indiscriminately together with other NPN containing feeds. Consult an AFGRI Animal Feeds Technical Advisor.
- Protect the feed against rain. Urea is water soluble and animals drinking such a solution could be poisoned.
- Vinegar is an effective remedy against NPN poisoning. Treatment: Mix with an equal amount of water and dose half a bottle per calf or large sheep and 2 to 4 bottles per head of cattle (1 bottle = 750ml).



12 Byls Bridge Boulevard Highveld Ext 73 Centurion

Tel: +27 11 063 2347 afgrianimalfeeds.co.za